

Spacetime at the Planck scale and Path Integral

G. Mangano

INFN, Sezione di Napoli, Mostra d'Oltremare Pad. 20, I-80125 Napoli, Italy

E-mail: mangano@na.infn.it

We discuss a path integral formalism to introduce noncommutative generalizations of spacetime manifold in even dimensions, which have been suggested to be reasonable effective pictures at very small length scales, of the order of Planck length.

*Amor, ch'a nullo amato amar perdona,
mi prese del costui piacer sì forte,
che, come vedi, ancor non m'abbandona.*

To Carmen

I. INTRODUCTION

A fascinating open problem of modern physics is a satisfactory understanding of the structure of spacetime at very short length scale, of the order of the Planck length $\lambda_P \sim 10^{-33}$ cm. On one side the difficulties of quantization of gravity in canonical ways can be only due to technical reasons, and we have simply to wait for new powerful mathematical tools. What is implicit in this possibility is however that the very basic structure of spacetime is a customary manifold with suitable differential and topological structures, which is the arena for interactions among quantized fields. On the other hand, many arguments on operational limits on position and time measurements have been considered in literature [1], suggesting that this description should be modified in a way which is reminiscent of the quantization of phase space in ordinary quantum mechanics. It is worth stressing that, in this case, it is the very notion of *manifold* which now undergoes a dramatic change. If we adopt a dual point of view, it is well known that the (commutative) algebra of smooth functions over spacetime manifold contains, using Gelf'and–Naimark reconstruction theorem, all informations on the underlying space. Switching on the Planck length then amounts to consider the noncommutative algebra still generated by positions and time X^μ , now looked upon as generators with nontrivial commutation relations

$$[X^\mu, X^\nu] = i\lambda_P^2 Q^{\mu\nu}, \quad (1.1)$$

where the antisymmetric tensor $Q^{\mu\nu}$, depends, in general, on the X^μ . The noncommutative $*$ -algebra $\mathcal{A}$, generated by regular representations of Eq. (1.1), contains all informations on what we may call *noncommutative spacetime*, while the geometric picture, which greatly helps in the commutative cases, is lost.

The analogy with ordinary quantum mechanics and quantization of phase space strongly suggests to those who are fascinated by the path integral approach, as the author of the present paper is, to establish a path integral formulation of these noncommutative geometries, taking the point of view that a class of linear functionals over $\mathcal{A}$, which turn into evaluation maps in the commutative limit, can be expressed as integrals, with a suitable measure, of ordinary functions over the classical, commutative spacetime. The reason why such an approach could eventually turn out to be useful is that it relates in a rather simple way the algebra generated by the X^μ to its commutative counterpart, and this is a powerful way to look for generalization at the noncommutative level of the basic structures of Riemannian geometry: metric, connections and curvature.

II. PATH INTEGRAL OVER SPACETIME

We start considering the pair (M^{2n}, Ω) , where M^{2n} is an even-dimensional differentiable manifold and Ω a symplectic form, and introduce the *generating functional*

$$Z(x_0, J) = N \int D[\gamma] \exp \left[i\lambda_P^{-2} \left(\int_\Gamma \Omega + (x, J) \right) \right], \quad (2.1)$$

where N is a normalization constant, J an arbitrary source, the integration is carried over all closed curves γ with base point $x_0 \in M^{2n}$ and Γ is any two-dimensional surface with boundary γ . We have also defined

$$(x, J) = \int_{\gamma} x^{\mu}(\tau) J_{\mu}(\tau) d\tau, \quad (2.2)$$

where $x^{\mu}(\tau)$ is any parameterization of γ . We here illustrate the basic construction for a manifold with trivial second homology group. The general case will be discussed in section 3. We therefore have

$$\int_{\Gamma} \Omega = \int_{\gamma} A_{\mu}(x) \frac{d}{d\tau} x^{\mu}(\tau) d\tau, \quad (2.3)$$

with $\Omega = dA$. $Z(x_0, J)$ is invariant under curve reparameterization, provided we redefine the arbitrary external source J . In the following we will consider $\tau \in [0, 1]$. Equation (2.1) is clearly inspired by the expression of path integral in phase space with external sources in the coherent state representation. The generating functional $Z(x_0, J)$ defines a (noncommutative) algebra $\mathcal{A}$ generated by $2n$ elements X^{μ} , implicitly defined via the introduction of a set of linear functionals ρ_{x_0} as follows

$$\begin{aligned} \rho_{x_0}(X^{\mu_1} \dots X^{\mu_k}) &\equiv \lim_{\tau_i - \tau_{i-1} \rightarrow 0^-} N Z(x_0, 0)^{-1} \int D[\gamma] e^{i \int_{\Gamma} \Omega / \lambda_P^2} x^{\mu_1}(\tau_1) \dots x^{\mu_k}(\tau_k) \\ &= \lim_{\tau_i - \tau_{i-1} \rightarrow 0^-} (-i \lambda_P^2)^k Z(x_0, 0)^{-1} \frac{\delta}{\delta J_{\mu_1}(\tau_1)} \dots \frac{\delta}{\delta J_{\mu_k}(\tau_k)} Z(x_0, J) \Big|_{J=0}. \end{aligned} \quad (2.4)$$

The way the limit is taken guarantees the ordering of the X^{μ_i} in the right-hand side of Eq. (2.4). Once defined on all polynomials in the X^{μ} , Eq. (2.4) can be applied to the entire algebra of continuous functions in the weak topology.

To understand this definition, we consider the *classical* limit $\lambda_P \rightarrow 0$. The integral over curves is then dominated by the contribution at stationary points, i.e. $\Omega_{\mu\nu}(x) dx^{\nu} / d\tau = 0$. Since Ω is not degenerate, the leading term satisfying $x^{\mu}(0) = x^{\mu}(1) = x_0^{\mu}$, is therefore given by the curve $x^{\mu}(\tau) = x_0^{\mu}$, $\forall \tau$, and thus

$$\rho_{x_0}(X^{\mu_1} \dots X^{\mu_k}) \rightarrow x_0^{\mu_1} \dots x_0^{\mu_k} + \mathcal{O}(\lambda_P^2). \quad (2.5)$$

The maps ρ_{x_0} reduce to the evaluation maps ($*$ -homomorphisms) of the commutative algebra of smooth functions over M^{2n} , $\rho_{x_0} : f(x) \mapsto f(x_0)$

The explicit evaluation of $Z(x_0, J)$ may be rather involved or even impossible for a generic two form Ω ; in these cases application of perturbation techniques, as in ordinary path integral in quantum mechanics, may nevertheless provide some information on the underlying noncommutative spacetime. In the simple case of a constant Ω , or assuming that it is a slowly varying function of x on the scale λ_P , one can easily compute Eq. (2.1) finding [2]

$$\rho_{x_0}(X^{\mu}) = x_0^{\mu}, \quad \rho_{x_0}([X^{\mu}, X^{\nu}]) = \frac{i}{2} \lambda_P^2 \Omega^{-1\mu\nu}(x_0). \quad (2.6)$$

As expected the commutator is proportional to the inverse symplectic form Ω^{-1} and using notation of Eq. (1.1), we see that X^{μ} generates a noncommutative algebra with $\rho_{x_0}(Q^{\mu\nu}) = \Omega^{-1\mu\nu}(x_0)/2$.

III. NON-TRIVIAL TOPOLOGIES AND BLACK HOLE AREA QUANTIZATION

In the case of a manifold M^{2n} with nontrivial second homology group the measure in the integral over closed curves in Eq. (2.1) is, in general, a multivalued function, since the integral of Ω over two surfaces Γ and Γ' , both with boundary γ , may be different if $\Gamma - \Gamma'$ is a non trivial 2-cycle. The ambiguity in the choice of the surface Γ , however, can be removed by requiring *quantization* conditions on the integral of Ω over a set of generators Γ_i of $H^2(M^{2n})$ such that $\exp(i \int_{\Gamma_i} \Omega \lambda_P^{-2})$ is single valued, analogous to the one introduced to quantize the motion of a particle in the field of a magnetic charge.

Let us consider for example the pair (S^2, Ω) . The generalization to an arbitrary manifold is straightforward [2]. For any closed curve the measure in the path integral will be always single valued if we require that the integral of Ω over S^2 is a multiple of 2π , in unit λ_P^2

$$\int_{S^2} \Omega = 2n\pi \lambda_P^2 \quad (3.1)$$

We stress the point that requiring Eq. (3.1) is a consistency condition to construct, via the path integral approach outlined in this paper, a deformation of spacetime, and so it seems to us intimately related to the appearance of

microscopic noncommutativity at the Planck scale. Actually, we observe that what we have just discussed could have an intriguing relationship with the idea that black hole horizon area is quantized and its spectrum is uniformly spaced [3]. This is in fact what condition (3.1) is stating, if S^2 represents the black hole horizon area and with Ω the area two-form. Black hole physics is probably one of the best scenarios where the structure of spacetime at small scales, otherwise unobservable, could be felt well above the Planck length. The topological constraint (3.1) may be, perhaps, a clue in the direction of a deep interplay of microscopic noncommutativity and macroscopic phenomena such as black hole mass quantization.

-
- [1] D. Amati, M. Ciafaloni and G. Veneziano, *Nucl. Phys.* **B347**, 550 (1990);
 S. Doplicher, K. Fredenhagen and J. E. Roberts, *Comm. Math. Phys.* **172**, 187 (1995);
 H. S. Snyder, *Phys. Rev.* **71**, 38 (1947);
 A. Kempf, G. Mangano and R. B. Mann, *Phys Rev.* **D52**, 1108 (1995); A. Kempf and G. Mangano, *Phys. Rev.* **D55**, 7907 (1997), and references therein.
 - [2] G. Mangano, *J. Math. Phys.* **39**, 2584 (1998).
 - [3] For a review see J. D. Bekenstein, gr-qc/9710076.